

A review of camera trapping for conservation behaviour research

Caravaggi, A.; Banks, P.B.; Burton, A.C. ; Finlay, C.M.V.; Haswell, Peter;
Hayward, Matthew; Rowcliffe, J.M.; Wood, M.D.

Remote Sensing in Ecology and Conservation

DOI:
[/10.1002/rse2.48](https://doi.org/10.1002/rse2.48)

Published: 01/09/2017

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):

Caravaggi, A., Banks, P. B., Burton, A. C., Finlay, C. M. V., Haswell, P., Hayward, M., Rowcliffe, J. M., & Wood, M. D. (2017). A review of camera trapping for conservation behaviour research. *Remote Sensing in Ecology and Conservation*, 3(3), 109-122. <https://doi.org/10.1002/rse2.48>

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Review

A review of camera trapping for conservation behaviour research

Caravaggi, A.¹, Banks, P.B.², Burton, A.C.³, Finlay, C.M.V.⁴, Haswell, P.M.⁵, Hayward, M.W.^{5,6}, Rowcliffe, J.M.⁷, Wood, M.D.⁸

¹ School of Biological Sciences, Queen's University Belfast, Belfast, BT9 7BL, UK

² School of Life and Environmental Sciences, The University of Sydney, NSW 2006

³ Department of Forest Resources Management, University of British Columbia, 2215 - 2424 Main Mall, Vancouver, BC, Canada, V6T 1Z4

⁴ Ulster Wildlife, McClelland House, 10 Heron Road, Belfast, BT3 9LE

⁵ School of Biological Sciences, Bangor University, Bangor, Gwynedd, LL57 2UW, UK.

⁶ Centre for African Wildlife Ecology, Nelson Mandela University, Port Elizabeth 6031 South Africa; and Centre for African Wildlife Ecology, University of Pretoria, Pretoria, 0001 South Africa

⁷ ZSL Institute of Zoology, Regent's Park, London NW 4RY, UK

⁸ School of Environment & Life Sciences, University of Salford, Manchester, M5 4WT, UK

Corresponding author email: acaravaggi01@qub.ac.uk

Running title: Camera traps and conservation behaviour

Word count

Abstract:	211
Main text:	4,297
Acknowledgments:	95
References:	3,678
Tables:	556
Figures:	113

Tables: 2

Figures: 1

Abstract

An understanding of animal behaviour is important if conservation initiatives are to be effective. However, quantifying the behaviour of wild animals presents significant challenges. Remote-sensing camera traps are becoming increasingly popular survey instruments that have been used to non-invasively study a variety of animal behaviours, yielding key insights into behavioural repertoires. They are well-suited to ethological studies and provide considerable opportunities for generating conservation-relevant behavioural data if novel and robust methodological and analytical solutions can be developed. This paper reviews the current state of camera-trap-based ethological studies, describes new and emerging directions in camera-based conservation behaviour, and highlights a number of limitations and considerations of particular relevance for camera-based studies. Three promising areas of study are discussed: i) documenting anthropogenic impacts on behaviour; ii) incorporating behavioural responses into management planning; and iii) using behavioural indicators such as giving up densities and daily activity patterns. We emphasise the importance of reporting methodological details, utilising emerging camera trap metadata standards and central data repositories for facilitating reproducibility, comparison and synthesis across studies. Behavioural studies using camera traps are in their infancy; the full potential of the technology is as yet unrealised. Researchers are encouraged to embrace conservation-driven hypotheses in order to meet future challenges and improve the efficacy of conservation and management processes.

Key words: Ethology, remote sensing, anthropogenic impacts, behavioural indicators, monitoring, management

58 **Introduction**

59 Animal behaviour is an important component of conservation biology (Berger-Tal et al. 2011),
60 and, hence, is of considerable interest to researchers and wildlife managers (Caro and Durant
61 1995). For example, behavioural studies can increase our understanding of species' habitat
62 requirements (Pienkowski 1979), reproductive behaviour (Cant 2000) and dispersal or
63 migration (Doerr et al. 2011), and elucidate impacts of habitat fragmentation (Merckx and Van
64 Dyck 2007) or climate change (Moller 2004). Animal behaviour can also be a useful
65 monitoring tool, with individual and group-level responses used to evaluate the impacts of
66 management (Morehouse et al. 2016). It is important, therefore, to incorporate behaviour into
67 conservation planning; its omission limits efficacy of conservation actions and could lead to
68 failure (Berger-Tal et al. 2011). The confluence of conservation biology and ethology has come
69 to be known as 'conservation behaviour', wherein conservation problems are addressed by the
70 application of behavioural research (Blumstein and Fernández-Juricic 2004; Berger-Tal et al.
71 2011).

72 Quantifying the behaviour of wild animals presents significant challenges. Direct
73 observation of animals can allow the evaluation of individual responses to environmental
74 stimuli. Such studies may be weakened, however, by the influence of the human observer on
75 focal animals (Nowak et al. 2014) and limited by small sample size and logistical constraints
76 (Bridges and Noss 2011). Furthermore, only a limited number of species and habitats are
77 amenable to direct, field-based observations (e.g. larger species and those that can be
78 habituated; and in open and accessible habitats). Many of these have already been the focus of
79 direct behavioural research (Schaller 1967; Kruuk 1972; Caro 1994) or may be atypical of more
80 common habitats and can lead to inconsistent results (Laurenson 1994 vs Mills and Mills 2014).
81 In cases where focal animal(s) cannot easily be directly observed, the vast majority of field-
82 based behavioural studies have used radio (VHF) or satellite (GPS) telemetry, activity sensors

and/or biologgers (e.g. Lewis et al. 2002; Grignolio et al. 2004; Shamoun-Baranes et al. 2012; Bouten et al. 2013). The advantages and disadvantages of these methods, which are currently the gold-standards for obtaining spatio-temporal behavioural data, are summarised in Table 1, highlighting that while these devices can provide powerful insights, they also have significant logistical and inferential limitations. Consequently, the suite of species that have had their behaviour quantified is biased and limited. New methods of obtaining behavioural data are, therefore, urgently required.

Camera traps (i.e. cameras that are remotely activated via an active or passive sensor; hereafter referred to as CTs) offer a reliable, minimally-invasive, visual means of surveying wildlife that substantially reduces survey effort. CTs are increasingly popular in ecological studies (Burton et al. 2015; Rovero and Zimmermann 2016) and provide a wealth of information that is often of considerable conservation value (e.g. Ng et al. 2004; Di Bitetti et al. 2006; Caravaggi et al. 2016). Continued technological improvements and decreasing equipment costs (Tobler et al. 2008a), combined with their demonstrated versatility (Rovero et al. 2013), mean that CTs will only continue to grow in popularity. CT data take the form of a still image or video of an individual or a group of individuals, of one or more species, which have been detected within the camera and location-specific zone of detection. These images can be linked with additional information, including the date, time and location at which the image was recorded. CT surveys have been effectively used to quantify species diversity (Tobler et al. 2008b), relative abundance (Carbone et al. 2001; Villette et al. 2017) and population parameters (Karanth et al. 2006; Rowcliffe et al. 2008), demonstrate site occupancy of rare or cryptic species (Linkie et al. 2007) and describe species replacement processes (Caravaggi et al. 2016). CTs have also been used in behavioural studies (Bridges and Noss 2011; Maffei et al. 2005). In a recent review of 266 CT studies, Burton et al. (2015)

characterized one-third as addressing behavioural questions (e.g. activity patterns, diet; Table 2).

In this paper, we review some of the recent literature on animal behaviour as elucidated by camera trapping studies. We then describe a number of common issues encountered by researchers undertaking such surveys and, finally, suggest future avenues of research that may be of considerable benefit to conservation initiatives. This review serves as a point of reference for researchers and practitioners undertaking conservation-oriented CT surveys of animal behaviour.

Current applications of camera traps to animal behaviour

CTs are well-suited to ethological studies, providing increasing opportunities to undertake extensive and detailed sampling of wild animal behavioural repertoires (see Fig. 1 and Table 2 for examples). The nature of the technology confers a number of important benefits. For example, CTs facilitate detailed studies of behaviours in species that were previously considered too small or elusive to be reliably observed in the field. CTs have been used to understand burrowing behaviour in <40g northern hopping-mice (*Notomys aquilo*; Diete et al. 2014) and olfactory communication in native and introduced <120g rats (*Rattus* sp.; Heavener et al. 2014). The use of CTs may also lead to a reduction in observer bias as, while a human observer is required to review collected images and assign individual and/or species identities and behaviours, cameras allow independent verification and recurrent analysis of observations. This is in contrast to conventional field methods for documenting behaviour, where it is rarely possible for another scientist to independently verify observational data.

Many types of animal behaviours have been studied with CTs (Table 2), including foraging (Otani 2001), daily activity patterns (Tan et al. 2013), scent marking (Delgado-V. et al. 2011), movement (Ford et al. 2009), livestock depredation (Bauer et al. 2005), and use of a variety of habitat features including dens/burrows (Clapham et al. 2014), urban habitats (Marks and Duncan 2009), corridors (LaPoint et al. 2013) and waterholes (Hayward and Hayward 2012). CT studies have often yielded key behavioural insights that may otherwise have remained unknown, many of which could be important to conservation processes. For example, studies investigating the efficacy of highway crossings in Banff National Park, Canada, described the effectiveness of under- and over-passes, an expensive and controversial means of impact mitigation (Clevenger and Waltho 2000; Ford et al. 2009), which is now being duplicated in other parts of the world. Picman and Schriml (1994) observed the predators of quail (*Coturnix coturnix*) nests in a variety of habitats, elucidating temporal variation and relative importance of each predatory species. The application of this method to the study of threatened avifauna has clear conservation benefits via the identification of direct impacts on egg success and the development of appropriate mitigation and monitoring techniques. Similarly, cameras provide more accurate post-hibernation den-emergence estimates for American black bears (*Ursus americanus*) than conventional methods, i.e. den visits and radio telemetry (Bridges et al. 2004). Long-term monitoring of emergence relative to climate may yield important insights into the effects of climate change on black bears and other hibernating species (*sensu* Bridges and Noss 2011).

The majority of ethological CT studies conducted thus far have been primarily curiosity-driven, rather than being motivated by applied conservation-focussed hypotheses. This is not to say that a large number of these studies do not have conservation value. On the contrary, the conservation relevance of the data is often explicitly discussed. It is apparent, however, that there is an increasing need for conservation-driven studies. CTs are among the

most promising and flexible tools available and we are only beginning to explore their potential.

Emerging directions in camera-based conservation behaviour

The growth in popularity and application of CT surveys and novel solutions to non-behavioural questions of animal ecology (e.g. Rowcliffe et al. 2008; Martin et al. 2015; Bowler et al. 2016) suggests that creative methodological and analytical solutions will be increasingly used to investigate animal behaviours. If these novel studies are to be developed, it is important that researchers strive for true experimental designs focussed on conservation behaviour. A particular strength of CT surveys is the potential for multiple studies to be carried out concurrently (e.g. estimation of focal species population density and the species richness of the surveyed area). Thus, behaviour can be recorded alongside other important parameters, thereby facilitating insight into processes such as density-dependent behaviours and responses to climate change. New approaches are also being developed to move beyond correlational approaches and incorporate CTs into manipulative experiments, such as measuring animal behavioural responses to introduced stimuli (e.g. predator calls; Suraci et al. 2016).

Berger-Tal et al. (2011) described three ways in which behavioural research can be of conservation benefit: i) identifying the impact of anthropogenic environmental changes on behaviour; ii) considering behavioural aspects of conservation initiatives ('behaviour-based management'); and iii) identifying behavioural indicators which are suggestive of changes in populations or the environment. We use this framework as a basis for our recommendations, below.

Anthropogenic impacts

179

180 An important area of conservation research lies in understanding the influence of
181 anthropogenic stressors on animal behaviours and predicting the resulting population-level
182 responses in order to inform management practices. Stressors such as habitat fragmentation,
183 disturbance, the creation of ecological traps and the introduction of non-native species can have
184 significant effects on behaviour (Robertson and Hutto 2006) and, hence, fitness (Berger-Tal et
185 al. 2011). For example, animals may exhibit increasing wariness in areas of greater disturbance
186 (Stewart et al. 2016) and may change their daily activity patterns in close proximity to human
187 populations (Carter et al. 2012). While anthropogenic impacts are generally negative, some
188 species show benefits such as increased occupancy in fragmented landscapes (Fleschutz et al.
189 2016), or using human activity to evade apex predators (Muhly et al. 2011; Steyaert et al. 2016).
190 Impacts on one species may also have spillover effects on the wider ecological community
191 (Wright et al. 2010; Clinchy et al. 2016).

192 Habitat fragmentation, the division of large, connected habitats into small, isolated
193 fragments separated by dissimilar habitats, is a major conservation issue (Haddad et al. 2015).
194 Fragmentation has a wide range of potential impacts on species and ecosystems (e.g. via edge
195 effects, patch size, shape and complexity and distance from other patches; Fahrig 2003), and
196 these impacts may be mediated through effects on animal behaviour. CTs provide new
197 opportunities for documenting behavioural responses to fragmentation. For example, the
198 activity patterns of nine-banded armadillos (*Dasypus novemcinctus*) varied in association with
199 forest patch size, among other factors, while patch time-since-isolation was predictive of agouti
200 (*Dasyprocta leporina*) activity (Norris et al. 2010).

201 The disruption of dispersal behaviour can lead to the endangerment and potential
202 extinction of isolated populations by various mechanisms, including changes to genetic
203 diversity and structure (Keyghobadi 2007), stochastic threats (Fischer and Lindenmayer 2007)

and long-term displacement effects (Ewers and Didham 2005). Using CTs to document dispersal behaviour can improve understanding of responses to movement disruption (Blumstein and Fernández-Juricic 2004) and inform design and implementation of mitigation measures that encourage dispersal. Aimed at species with individually-identifiable markings or tags, individual-level analysis is potentially possible, although inferences about dispersal can also be drawn without individual identification. For examples, cameras are well suited to quantifying use of presumed dispersal routes or movement corridors, including mitigations designed to promote connectivity (e.g. highway crossings; Clevenger and Waltho 2005; Ford et al. 2009). CTs can also be used to identify colonization of new habitat patches (including range expansions or species invasions) and parameterize landscape connectivity models (Brodie et al. 2015).

No studies have integrated environmental sensors into CT studies investigating anthropogenic impacts on behaviour, and we believe this is a promising area for future development. Local temperature, precipitation and humidity can readily be recorded, and phenocams can be used to document vegetation and environmental changes (Brown et al. 2016). Collecting such information alongside CT-based behavioural data will allow us to increase our understanding of how animals respond to changing conditions at both large (population) and small (localities within home ranges) spatial scales. This is particularly important given the rapid changes that are predicted to occur under climate change.

Behaviour-based management

Berger-Tal et al. (2011) suggested that behaviour-sensitive management and behavioural modification are two key pathways through which ethology can inform active management for conservation. The former considers animal behaviour in the design of reserves and corridors,

228 planning species reintroductions and translocations, and epidemiology with the goal of
229 stabilising or increasing threatened populations or controlling pest or invasive species.
230 Behavioural modification focuses on changing or preserving key behaviours within a focal
231 population. CT surveys have the potential to inform both of these areas.

232 Considering social dynamics is one important area in which CT surveys can inform
233 behaviour-sensitive management. Social species, i.e. those that interact and/or live together,
234 often exhibit complex inter-group relationships and social structure (Rowell 1966; Creel 1997;
235 Archie et al. 2006; Wolf et al. 2007; Wey et al. 2008), that are susceptible to rapid change via
236 the social displacement or death of one or more individuals. This can have severe consequences
237 for the species and/or their environment (e.g. Nyakaana et al. 2001). Social Network Analysis
238 (SNA) facilitates the study of relationships between nodes (i.e. individuals), within networks
239 (i.e. social groups; Sueur et al. 2011). The methodology is increasingly used to study animal
240 behaviour (Lusseau et al. 2006; Whitehead 2008; Voelkl and Kasper 2009; Jacoby and
241 Freeman 2016). Examples of SNA demonstrating a direct benefit to conservation, however, are
242 few. SNA studies are limited in that they require the reliable identification of individuals and,
243 hence, are only applicable with CTs where animals exhibit individual characteristics or
244 markings, or where marks (e.g. tags) can be attached. However, placing cameras in areas
245 frequented by social groups such as feeding or resting sites, and with a sufficient number of
246 units, could yield a considerable amount of important data for behaviour-sensitive
247 management. Such site-specific studies have some limitations and incur biases that require
248 evaluation. For example, individuals may not be equally detectable, or full groups may not be
249 observed. Furthermore, it would be difficult to account for behaviours and social interactions
250 which occur while away from the focal site. However, SNA analyses do not require constant
251 observation of all group members to be effective (see Jacoby and Freeman 2016). Assessing

potential bias with calibration by direct observation or other methods and placing observations in appropriate contexts is therefore important.

SNA has the potential to increase our understanding of disease or pathogen transmission and individual or group vulnerability (Krause et al. 2007), an issue of particular relevance to the conservation of species which are susceptible to outbreaks (e.g. Hamede et al. 2009). SNA studies have demonstrated that the removal of certain individuals (e.g. via hunting) can have a considerable effect on the stability of the social network (e.g. Flack et al. 2006), thus demonstrating their potential utility in elucidating the impacts of the bushmeat trade on inter- and intra-group dynamics in primates, for example. Furthermore, SNA has implications for reintroduction programmes, where the (re)construction of cohesive social structures in a captive setting would be necessary for the return of the focal species to the wild (Abell et al. 2013). Studies of the relationships between individuals, therefore, can help us to understand how social behaviour is influenced by a variety of factors and, hence, provide an additional means by which practitioners can build an evidence base to address conservation questions.

CTs can also be applied to studies of behavioural modification. For example, Davies et al. (2016) investigated responses of African herbivores to changes in predation risk resulting from recently-reintroduced lions. Cameras could also be used to monitor animal responses to conflict mitigation measures such as the use of bees or chilli to deter crop-raiding elephants (Karidozo and Osborn 2015; Ngama et al. 2016).

Behavioural indicators

The ways in which animals adapt their foraging behaviour in human-impacted environments have important implications for their abilities to adapt and persist under increasing pressures. Behavioural indicators can be used to assess the state of animals and the environments they

inhabit, highlighting important conservation issues such as population decline or habitat degradation, or being used to monitor the efficacy of management (Berger-Tal et al. 2011). Behaviour effectively acts as an early-warning system, indicating changes to processes before they are evident through, for example, population decline.

The giving up density (GUD; i.e. the amount of food left behind from a known starting quantity; Brown 1988) is one such behavioural indicator that has been used to study predation risk (Orrock 2004; Severud et al. 2011), energetic costs (Nolet et al. 2006), forager state and forage quality (Hayward et al. 2015), plant toxins (Emerson and Brown 2015), competition (Brown et al. 1997) and predator-prey dynamics (Andruskiw et al. 2008). It is also central to describing the “landscape of fear” (i.e. relative levels of predation risk within an area of use) of an animal and its habitat preferences, which are direct behavioural indicators with significant conservation implications (Kotler et al. 2016). CTs offer a relatively reliable way of using the GUD technique to ask more in-depth questions of conservation relevance. For example, CTs have been used to calculate GUDs for multiple species (Lerman et al. 2012), examine (Mella et al. 2015), and differentiate individual versus group foraging habits (Carthey and Banks 2015). These observations can then be used to inform the development of hypotheses relating to the broader effects of local food and predator abundance, predation pressure and inter- and intra-specific competition. With advancements in CT technology and creative experimental design, a wealth of conservation-focussed GUD applications are now possible.

A key strength of CTs lies in collecting data on multiple species, either as bycatch in a focal study, or as part of a specific multi-taxa investigation. Accordingly, there has been an increasing focus on assessing species interactions and niche partitioning via comparisons of co-occurrence and activity patterns (de Almeida Jacomo et al. 2004; Kukielka et al. 2013; Farris et al. 2014; Wang et al. 2015; Bu et al. 2016; Cusack et al. 2016; Sweitzer and Furnas 2016). Animal activity patterns are shaped by a number of factors, including foraging

efficiency (Lode 1995), predator/prey activity (Middleton et al. 2013), photoperiodism (McElhinny et al. 1997), and competition (Rychlik 2005). Conservation-focussed studies using these methodologies, however, are scarce. Changes in the way species interact and use the landscape may be indicative of responses to changing environmental pressures and, hence, can direct development of early conservation strategies. For example, brown bears (*Ursus arctos*; Ordiz et al. 2013) altered their movement patterns and wolverines (*Gulo gulo*; Stewart et al. 2016) behaved differently when faced with human disturbance, potentially impacting their ecosystem roles and, hence, associated species and habitats. Disturbance of the activity patterns of one or more species in a dynamic interaction, particularly ecological competitors or predators and prey, can therefore be interpreted as indicative of environmental changes and, hence, suggest additional lines of enquiry and highlight areas of conservation concern.

Scaling-up

Cameras can be used to monitor large-scale biodiversity conservation processes (O'Brien et al. 2010; Ahumada et al. 2013) and investigate animal behaviour on a landscape scale. Scaling-up CT networks would provide stronger, larger-scale inferences on spatio-temporal variation in behaviours (Steenweg et al. 2016). Studies conducted on a broader scale have inherent limitations, however, that are not necessarily considerations for more localised investigations. The trade-off between the scale of investigation and camera array density has spatio-temporal implications which must be considered when designing a study, formulating hypotheses and deriving inferences from resultant data. Broad-scale studies are also ostensibly limited by the number of researchers available to place and check cameras and process data. The recruitment of volunteers (i.e. citizen scientists), however, offers a means of expanding the scope of research (Cohn 2008), greatly expanding spatial coverage and delivering a wealth of temporally

comparable data (McShea et al. 2016). Emerging large-scale camera monitoring initiatives, such as Snapshot Serengeti (www.snapshotserengeti.org; Swanson et al. 2015) and Wildcam Gorongosa (www.wildcamgorongosa.org) demonstrate the benefits of this approach. CT projects utilising citizen science have the potential to deliver a substantial amount of behavioural data (McShea et al. 2016) and inform conservation processes. However, few large-scale studies utilising citizen science involve behavioural analyses. CT video data can produce vast amounts of video footage but the extraction of key behavioural data from video footage is time consuming, imposing a major obstacle. Crowdsourcing video interpretations can overcome this limitation, however, and the use of robust ethograms, simple training regimes and blinding of observers to treatments can assuage concerns about the reliability of citizen science interpretations (e.g. Carthey 2013).

Synthesising across projects offers another means of conducting broader analyses (Steenweg et al. 2016). We recommend that researchers embrace emerging CT metadata standards and associated opportunities to use common data repositories such as Wildlife Insights (www.wildlifeinsights.org; Forrester et al. 2016), thus increasing the potential for the synthesis of inferences across large scales. The value of current data repositories is reduced, however, by their reliance on static images and omission of video. While it is possible to derive important behavioural data from still images, videos are undoubtedly more informative and an important future direction for CT-based behavioural research. Expenses notwithstanding, it is in the interests of conservation behaviour researchers to establish a digital repository for video data.

Relevant limitations and considerations

Despite the great promise of new insights in conservation behaviour from CTs, it is important to consider potential limitations. CTs are passive instruments; thus, while it is possible to identify animals according to species, age-class (Clapham et al. 2014), sex (Bezerra et al. 2014) or, indeed, identify individuals (Karanth et al. 2006; Zheng et al. 2016), the collection of biometric, genetic and other data of interest requires the application of supplementary or alternative methodologies. Furthermore, CTs are frequently considered to be non-intrusive, causing little to no disturbance. However, while the sound produced by recording units is largely inaudible to humans, it is frequently detected by wildlife (Meek et al. 2014a). Similarly, CTs which utilise visible light (as opposed to infra-red) increase the chances of the camera being detected by animals, potentially disrupting their natural behaviour (Meek et al. 2016a).

Camera failure, although rare, can result in the loss of large quantities of data. Similarly, camera theft is becoming increasingly common (Meek et al. 2016b). It is therefore necessary to balance the frequency of visits to maintain CTs with risk of data loss. To accommodate this, it is advisable to build some redundancy into the study design, such as the use of cameras that allow the transmission of images via Global Packet Radio Service (GRPS) and/or Wi-Fi and can therefore facilitate remote data collection and inform the timing of maintenance visits.

Detailed analysis of a target species' behavioural repertoire requires the use of video footage which often exposes the technical limitations of CT equipment. Many cameras offer only limited length of videos (e.g. 60 seconds), requiring the camera to be retriggered to continue the capture of the behaviours and, hence, creating gaps in the observation. Some cameras have a slow trigger time meaning that initial behaviours, which might be the most important in terms of measuring detection of a stimulus (rather than the response), can be missed. Sampling the behaviours of small species can be particularly challenging, with CTs typically designed for deer-sized game species (Weerakoon et al. 2014), a problem that will

require novel solutions. For example, flash-illuminated images are frequently obscured by overexposure when close enough to small mammals to observe behaviour clearly, whereas at the correctly exposed distance, animals can be too far away to reliably identify species or discern behaviours. Furthermore, understanding the reliability of camera surveys for addressing multi-species objectives remains an important area of methodological research (see Burton et al. 2015). Multi-taxa studies also require careful planning to ensure that CTs are appropriately located and adequately spaced to maximise the chances of capturing a diverse species assemblage. The choice and placement of cameras should, therefore, be dictated by the objectives of the study, the ecology of the study species, the statistical sampling framework and associated considerations.

An oft-repeated concern relates to study repeatability; specific details of study design (e.g. how survey sites were chosen, use of lures) and camera protocols (e.g. camera model, deployment details) are often lacking (Meek et al. 2014b; Burton et al. 2015). A number of factors influence the detection of individuals (see Burton et al. 2015) and sampling details may have important implications for analytical assumptions such as effective sampling area and site independence (Harmsen et al. 2010; McCoy et al. 2011; du Preez et al. 2014; Newey et al. 2015). Comprehensive methodological descriptions and utilisation of emerging CT metadata standards (Forrester et al. 2016) are important for facilitating reproduction, comparison and synthesis across studies.

Finally, as with any survey method, observations from CTs are incomplete and may contain biases that affect inferences. As noted above, species and individuals may vary in their detectability by CTs according to attributes such as body size, movement speed, curiosity and wariness. Behaviours observed by CTs may also not always be representative of behaviours more generally. It is thus incumbent upon researchers to remain vigilant for potential biases

and test CT-based inferences through comparison and calibration with more established ethological methods.

Conclusions

CTs are rapidly increasing in popularity, and their application to conservation behaviour is growing. Recent efforts to coordinate camera studies across large-scales through methodological standardization and/or better reporting of methodologies and metadata will facilitate broader ethological inferences on species' behavioural responses to environmental change. The development and application of new techniques and analytical methods explicitly focussed on anthropogenic impacts, behaviour-based management and behavioural indicators would undoubtedly benefit conservation programmes. CTs are not a panacea, but they confer many benefits to researchers and the diversity of possible applications is gradually being realised. We hope that this paper will act as a catalyst, advancing the adoption of CT technology within conservation behaviour. It is important, therefore, that potentially profitable avenues of investigation are identified and pursued if we are to maximise the generation of valuable data and, hence, improve the conservation outlook for the ever-increasing number of threatened or endangered species.

Acknowledgments

The authors thank the journal for their encouragement and support of this review and, specifically, the editor and reviewers for their valuable comments which strengthened the final manuscript. Authors are ordered alphabetically by surname, except for A. Caravaggi who

421 conceived, organised and compiled the manuscript. The contribution of M.D. Wood was
422 supported by the TREE project (www.ceh.ac.uk/TREE) funded by the Natural Environment
423 Research Council (NERC), Environment Agency and Radioactive Waste Management
424 Limited. The contributions of C.M.V. Finlay were supported by the Red Squirrels United
425 project (<http://www.redsquirrelsunited.org.uk/>) funded by EU Life and Heritage Lottery Fund.

426

References

- Abell, J. et al. 2013. A social network analysis of social cohesion in a constructed pride: implications for ex situ reintroduction of the African lion (*Panthera leo*). *PLoS ONE* 8(12), p. e82541.
- Ahumada, J.A. et al. 2013. Monitoring the status and trends of tropical forest terrestrial vertebrate communities from camera trap data: a tool for conservation. *PLoS ONE* 8(9), p. e73707.
- de Almeida Jacomo, A.T. et al. 2004. Niche separation between the maned wolf (*Chrysocyon brachyurus*), the crab-eating fox (*Dusicyon thous*) and the hoary fox (*Dusicyon vetulus*) in central Brazil. *Journal of Zoology* 262(1), pp. 99–106.
- Andruskiw, M. et al. 2008. Habitat-mediated variation in predation risk by the American marten. *Ecology* 89(8), pp. 2273–2280.
- Archie, E.A. et al. 2006. Dominance rank relationships among wild female African elephants, *Loxodonta africana*. *Animal Behaviour* 71(1), pp. 117–127.
- Azlan, J.M. and Sharma, D.S.K. 2006. The diversity and activity patterns of wild felids in a secondary forest in Peninsular Malaysia. *Oryx* 40(01), p. 36.
- Barron, D.G. et al. 2010. Meta-analysis of transmitter effects on avian behaviour and ecology: Meta-analysis of avian transmitter effects. *Methods in Ecology and Evolution* 1(2), pp. 180–187.
- Bauer, J.W. et al. 2005. Scavenging behaviour in puma. *The Southwestern Naturalist* 50(4), pp. 466–471.
- Berger-Tal, O. et al. 2011. Integrating animal behavior and conservation biology: a conceptual framework. *Behavioral Ecology* 22(2), pp. 236–239.
- Bezerra, B.M. et al. 2014. Camera trap observations of nonhabituated critically endangered wild blonde capuchins, *Sapajus flavius* (formerly *Cebus flavius*). *International Journal of Primatology* 35(5), pp. 895–907.
- Blumstein, D.T. and Fernández-Juricic, E. 2004. The emergence of conservation behavior. *Conservation Biology* 18(5), pp. 1175–1177.
- Bouten, W. et al. 2013. A flexible GPS tracking system for studying bird behaviour at multiple scales. *Journal of Ornithology* 154(2), pp. 571–580.
- Bowler, M.T. et al. 2016. Estimating mammalian species richness and occupancy in tropical forest canopies with arboreal camera traps. *Remote Sensing in Ecology and Conservation*. Available at: <http://doi.wiley.com/10.1002/rse2.35> [Accessed: 7 December 2016].
- Bridges, A.S. et al. 2004. Seasonal variation in American black bear *Ursus americanus* activity patterns: quantification via remote photography. *Wildlife Biology* 10(4), pp. 277–284.
- Bridges, A.S. and Noss, A.J. 2011. Behaviour and activity patterns. In: O’Connell, A. F. et al. eds. *Camera traps in animal ecology: Methods and analyses*. New York: Springer, pp. 57–69.

465 Brodie, J. et al. 2012. Climate change intensification of herbivore impacts on tree recruitment.
 466 *Proceedings of the Royal Society B: Biological Sciences* 279(1732), pp. 1366–1370.

467 Brodie, J.F. et al. 2015. Evaluating multispecies landscape connectivity in a threatened tropical
 468 mammal community: multispecies habitat corridors. *Conservation Biology* 29(1), pp. 122–132.

469 Brown, J.S. 1988. Patch use as an indicator of habitat preference, predation risk, and
 470 competition. *Behavioral Ecology and Sociobiology* 22(1), pp. 37–47.

471 Brown, J.S. et al. 1997. Competition between birds and mammals: a comparison of giving-up
 472 densities between crested larks and gerbils. *Evolutionary Ecology* 11(6), pp. 757–771.

473 Bu, H. et al. 2016. Spatial Co-Occurrence and Activity Patterns of Mesocarnivores in the
 474 Temperate Forests of Southwest China Allen, B. L. ed. *PLOS ONE* 11(10), p. e0164271.

475 Burton, A.C. et al. 2015. Wildlife camera trapping: a review and recommendations for linking
 476 surveys to ecological processes. *Journal of Applied Ecology* 52(3), pp. 675–685.

477 Cant, M.A. 2000. Social control of reproduction in banded mongooses. *Animal Behaviour*
 478 59(1), pp. 147–158.

479 Caravaggi, A. et al. 2016. An invasive-native mammalian species replacement process captured
 480 by camera trap survey random encounter models. *Remote Sensing in Ecology and Conservation*
 481 2(1), pp. 45–58.

482 Carbone, C. et al. 2001. The use of photographic rates to estimate densities of tigers and other
 483 cryptic mammals. *Animal conservation* 4(01), pp. 75–79.

484 Caro, T. and Durant, S.M. 1995. The importance of behavioral ecology for conservation
 485 biology: examples from Serengeti carnivores. In: Sinclair, A. R. E. and Arcese, P. eds.
 486 *Serengeti II: Dynamics, Management, and Conservation of an Ecosystem*. Chicago: University
 487 of Chicago, pp. 451–472.

488 Caro, T.M. 1994. *Cheetahs of the Serengeti Plains: group living in an asocial species*. Chicago:
 489 University of Chicago Press.

490 Carter, N.H. et al. 2012. Coexistence between wildlife and humans at fine spatial scales.
 491 *Proceedings of the National Academy of Sciences* 109(38), pp. 15360–15365.

492 Carthey, A.J.R. 2013. Naivete, novelty and native status: mismatched ecological interactions
 493 in the Australian environment. Australia: University of Sydney.

494 Carthey, A.J.R. and Banks, P.B. 2015. Foraging in groups affects giving-up densities: solo
 495 foragers quit sooner. *Oecologia* 178(3), pp. 707–713.

496 Carthey, A.J.R. and Banks, P.B. 2016. Naiveté is not forever: responses of a vulnerable native
 497 rodent to its long term alien predators. *Oikos* 125(7), pp. 918–926.

498 Clapham, M. et al. 2014. Scent-marking investment and motor patterns are affected by the age
 499 and sex of wild brown bears. *Animal Behaviour* 94, pp. 107–116.

500 Claridge, A.W. et al. 2004. Use of infrared digital cameras to investigate aspects of the social
501 behaviour of cryptic species. *Wildlife Research* 31(6), p. 645.

502 Clevenger, A.P. and Waltho, N. 2000. Factors influencing the effectiveness of wildlife
503 underpasses in Banff National Park, Alberta, Canada. *Conservation Biology* 14(1), pp. 47–56.

504 Clevenger, A.P. and Waltho, N. 2005. Performance indices to identify attributes of highway
505 crossing structures facilitating movement of large mammals. *Biological Conservation* 121(3),
506 pp. 453–464.

507 Clinchy, M. et al. 2016. Fear of the human ‘super predator’ far exceeds the fear of large
508 carnivores in a model mesocarnivore. *Behavioral Ecology*, p. arw117.

509 Cohn, J.P. 2008. Citizen science: can volunteers do real research? *BioScience* 58(3), p. 192.

510 Coulombe, M. et al. 2006. Quantification and accuracy of activity data measured with VHF
511 and GPS telemetry. *Wildlife Society Bulletin* 34, pp. 81–92.

512 Creel, S. 1997. Handling of African wild dogs and chronic stress: reply to East et al.
513 *Conservation Biology* 11(6), pp. 1454–1456.

514 Crook, N. et al. 2013. Bare-nosed wombats (*Vombatus ursinus*) use drainage culverts to cross
515 roads. *Australian Mammalogy* 35(1), p. 23.

516 Cusack, J.J. et al. 2016. Revealing kleptoparasitic and predatory tendencies in an African
517 mammal community using camera traps: a comparison of spatiotemporal approaches. *Oikos*.
518 Available at: <http://doi.wiley.com/10.1111/oik.03403> [Accessed: 18 December 2016].

519 Dalloz, M.F. et al. 2012. Positional behaviour and tail use by the bare-tailed woolly opossum
520 *Caluromys philander* (Didelphimorphia, Didelphidae). *Mammalian Biology - Zeitschrift für*
521 *Säugetierkunde* 77(5), pp. 307–313.

522 Davies, A.B. et al. 2016. Limited spatial response to direct predation risk by African herbivores
523 following predator reintroduction. *Ecology and Evolution* 6(16), pp. 5728–5748.

524 Delgado-V., C.A. et al. 2011. Behaviour of the Tayra *Eira barbara* near Medellín, Colombia:
525 preliminary data from a video-capturing survey. *Small Carnivore Conservation* 44(2011), pp.
526 19–21.

527 Di Bitetti, M.S. et al. 2006. Density, habitat use and activity patterns of ocelots (*Leopardus*
528 *pardalis*) in the Atlantic Forest of Misiones, Argentina. *Journal of Zoology* 270, pp. 153–163.

529 Diete, R.L. et al. 2014. Burrowing behaviour of the northern hopping-mouse (*Notomys aquilo*):
530 field observations. *Australian Mammalogy*. Available at:
531 <http://www.publish.csiro.au/?paper=AM13039> [Accessed: 26 March 2017].

532 Doerr, V.A.J. et al. 2011. Connectivity, dispersal behaviour and conservation under climate
533 change: a response to Hodgson et al.: Connectivity and dispersal behaviour. *Journal of Applied*
534 *Ecology* 48(1), pp. 143–147.

535 Edwards, S. et al. 2015. Spatiotemporal resource partitioning of water sources by African
536 carnivores on Namibian commercial farmlands: Carnivore resource partitioning at water
537 sources. *Journal of Zoology* 297(1), pp. 22–31.

538 Emerson, S.E. and Brown, J.S. 2015. The influence of food chemistry on food-safety tradeoffs
539 in samango monkeys. *Journal of Mammalogy* 96(1), pp. 237–244.

540 Emmons, L.H. et al. 2004. Bathing behavior of giant anteaters (*Myrmecophaga tridactyla*).
541 *Edentata* 6(1), p. 41.

542 Ewers, R.M. and Didham, R.K. 2005. Confounding factors in the detection of species responses
543 to habitat fragmentation. *Biological Reviews* 81(01), p. 117.

544 Fahrig, L. 2003. Effects of habitat fragmentation on biodiversity. *Annual Review of Ecology,*
545 *Evolution, and Systematics* 34(1), pp. 487–515.

546 Farris, Z.J. et al. 2014. Predator–Primate Distribution, Activity, and Co-occurrence in Relation
547 to Habitat and Human Activity Across Fragmented and Contiguous Forests in Northeastern
548 Madagascar. *International Journal of Primatology* 35(5), pp. 859–880.

549 Fischer, J. and Lindenmayer, D.B. 2007. Landscape modification and habitat fragmentation: a
550 synthesis. *Global Ecology and Biogeography* 16(3), pp. 265–280.

551 Flack, J.C. et al. 2006. Policing stabilizes construction of social niches in primates. *Nature*
552 439(7075), pp. 426–429.

553 Fleschutz, M.M. et al. 2016. Response of a small felid of conservation concern to habitat
554 fragmentation. *Biodiversity and Conservation* 25(8), pp. 1447–1463.

555 Ford, A.T. et al. 2009. Comparison of methods of monitoring wildlife crossing-structures on
556 highways. *Journal of Wildlife Management* 73(7), pp. 1213–1222.

557 Forrester, T. et al. 2016. An open standard for camera trap data. *Biodiversity Data Journal* 4,
558 p. e10197.

559 Grignolio, S. et al. 2004. Seasonal variations of spatial behaviour in female Alpine ibex (*Capra*
560 *ibex ibex*) in relation to climatic conditions and age. *Ethology Ecology & Evolution* 16(3), pp.
561 255–264.

562 Haddad, N.M. et al. 2015. Habitat fragmentation and its lasting impact on Earth’s ecosystems.
563 *Science Advances* 1(2), p. e1500052.

564 Hamede, R.K. et al. 2009. Contact networks in a wild Tasmanian devil (*Sarcophilus harrisii*)
565 population: using social network analysis to reveal seasonal variability in social behaviour and
566 its implications for transmission of devil facial tumour disease. *Ecology Letters* 12(11), pp.
567 1147–1157.

568 Harmsen, B.J. et al. 2010. Differential Use of Trails by Forest Mammals and the Implications
569 for Camera-Trap Studies: A Case Study from Belize: Trail Use by Neotropical Forest
570 Mammals. *Biotropica* 42(1), pp. 126–133.

571 Hayward, M.W. et al. 2015. Risk perception by endangered European bison *Bison bonasus* is
572 context (condition) dependent. *Landscape Ecology* 30(10), pp. 2079–2093.

573 Hayward, M.W. and Hayward, M.D. 2012. Waterhole use by African fauna. *South African*
574 *Journal of Wildlife Research* 42(2), pp. 117–127.

575 Heavener, S.J. et al. 2014. Competitive naïveté between a highly successful invader and a
576 functionally similar native species. *Oecologia* 175(1), pp. 73–84.

577 Jachowski, D.S. et al. 2015. Monitoring landscape-level distribution and migration phenology
578 of raptors using a volunteer camera-trap network. *Wildlife Society Bulletin* 39(3), pp. 553–563.

579 Jacoby, D.M.P. and Freeman, R. 2016. Emerging Network-Based Tools in Movement Ecology.
580 *Trends in Ecology & Evolution* 31(4), pp. 301–314.

581 Johnson, A. et al. 2006. Effects of human-carnivore conflict on tiger (*Panthera tigris*) and prey
582 populations in Lao PDR. *Animal Conservation* 9(4), pp. 421–430.

583 Karanth, K.U. et al. 2006. Assessing tiger population dynamics using photographic capture-
584 recapture sampling. *Ecology* 87(11), pp. 2925–2937.

585 Karidozo, M. and Osborn, F.V. 2015. Community based conflict mitigation trials: results of
586 field tests of chilli as an elephant deterrent. *Journal of Biodiversity & Endangered Species*
587 3(144). Available at: [http://esciencecentral.org/journals/community-based-conflict-mitigation-](http://esciencecentral.org/journals/community-based-conflict-mitigation-trials-results-of-field-tests-of-chilli-as-an-elephant-deterrent-2332-2543-1000144.php?aid=46217)
588 [trials-results-of-field-tests-of-chilli-as-an-elephant-deterrent-2332-2543-](http://esciencecentral.org/journals/community-based-conflict-mitigation-trials-results-of-field-tests-of-chilli-as-an-elephant-deterrent-2332-2543-1000144.php?aid=46217)
589 [1000144.php?aid=46217](http://esciencecentral.org/journals/community-based-conflict-mitigation-trials-results-of-field-tests-of-chilli-as-an-elephant-deterrent-2332-2543-1000144.php?aid=46217) [Accessed: 27 December 2016].

590 Keyghobadi, N. 2007. The genetic implications of habitat fragmentation for animals. *Canadian*
591 *Journal of Zoology* 85(10), pp. 1049–1064.

592 Kotler, B.P. et al. 2016. Direct behavioral indicators as a conservation and management tool.
593 In: Berger-Tal, O. and Saltz, D. eds. *Conservation Behavior*. Cambridge: Cambridge
594 University Press, pp. 307–351.

595 Krause, J. et al. 2007. Social network theory in the behavioural sciences: potential applications.
596 *Behavioral Ecology and Sociobiology* 62(1), pp. 15–27.

597 Kruuk, H. 1972. *The Spotted Hyaena*. Chicago: University of Chicago Press.

598 Kukiela, E. et al. 2013. Spatial and temporal interactions between livestock and wildlife in
599 South Central Spain assessed by camera traps. *Preventive Veterinary Medicine* 112(3–4), pp.
600 213–221.

601 LaPoint, S. et al. 2013. Animal behavior, cost-based corridor models, and real corridors.
602 *Landscape Ecology* 28(8), pp. 1615–1630.

603 Laurenson, M.K. 1994. High juvenile mortality in cheetahs (*Acinonyx jubatus*) and its
604 consequences for maternal care. *Journal of Zoology* 234(3), pp. 387–408.

605 Lerman, S.B. et al. 2012. Linking foraging decisions to residential yard bird composition. *PLoS*
606 *ONE* 7(8), p. e43497.

607 Leuchtenberger, C. et al. 2014. Activity patterns of giant otters recorded by telemetry and
608 camera traps. *Ethology Ecology & Evolution* 26(1), pp. 19–28.

609 Lewis, S. et al. 2002. Sex-specific foraging behaviour in a monomorphic seabird. *Proceedings*
610 *of the Royal Society of London B: Biological Sciences* 269(1501), pp. 1687–1693.

611 Linkie, M. et al. 2007. Estimating occupancy of a data deficient mammalian species living in
612 tropical rainforests: Sun bears in the Kerinci Seblat region, Sumatra. *Biological Conservation*
613 137(1), pp. 20–27.

614 Lode, T. 1995. Activity pattern of polecats *Mustela putorius* L. in relation to food habits and
615 prey activity. *Ethology* 100(4), pp. 295–308.

616 Lusseau, D. et al. 2006. Quantifying the influence of sociality on population structure in
617 bottlenose dolphins. *Journal of Animal Ecology* 75(1), pp. 14–24.

618 Maffei, L. et al. 2005. Ocelot (*Felis pardalis*) population densities, activity, and ranging
619 behaviour in the dry forests of eastern Bolivia: data from camera trapping. *Journal of Tropical*
620 *Ecology* 21(3), pp. 349–353.

621 Marks, B.K. and Duncan, R.S. 2009. Use of forest edges by free-ranging cats and dogs in an
622 urban forest fragment. *Southeastern Naturalist* 8(3), pp. 427–436.

623 Martin, E.H. et al. 2015. Modelling fine-scale habitat associations of medium-to-large forest
624 mammals in the Udzungwa Mountains of Tanzania using camera trapping. *Tropical Zoology*
625 28(4), pp. 137–151.

626 McCoy, J.C. et al. 2011. Bias associated with baited camera sites for assessing population
627 characteristics of deer. *The Journal of Wildlife Management* 75(2), pp. 472–477.

628 McElhinny, T.L. et al. 1997. Patterns of body temperature, activity, and reproductive behavior
629 in a tropical murid rodent, *Arvicanthis niloticus*. *Physiology & Behavior* 62(1), pp. 91–96.

630 McShea, W.J. et al. 2016. Volunteer-run cameras as distributed sensors for macrosystem
631 mammal research. *Landscape Ecology* 31(1), pp. 55–66.

632 Meek, P.D. et al. 2014. Camera traps can be heard and seen by animals. *PLoS ONE* 9(10), p.
633 e110832.

634 Meek, P. D. et al. 2014. Recommended guiding principles for reporting on camera trapping
635 research. *Biodiversity and Conservation* 23(9), pp. 2321–2343.

636 Meek, P.D. et al. 2016. Are we getting the full picture? Animal responses to camera traps and
637 implications for predator studies. *Ecology and Evolution* 6(10), pp. 3216–3225.

638 Meek, Paul D. et al. 2016. The higher you go the less you will know: placing camera traps high
639 to avoid theft will affect detection Rowcliffe, M. and De Angelo, C. eds. *Remote Sensing in*
640 *Ecology and Conservation* 2(4), pp. 204–211.

641 Mella, V.S.A. et al. 2015. Personality affects the foraging response of a mammalian herbivore
642 to the dual costs of food and fear. *Oecologia* 177(1), pp. 293–303.

643 Merckx, T. and Van Dyck, H. 2007. Habitat fragmentation affects habitat-finding ability of the
644 speckled wood butterfly, *Pararge aegeria* L. *Animal Behaviour* 74(4), pp. 1029–1037.

645 Middleton, A.D. et al. 2013. Linking anti-predator behaviour to prey demography reveals
646 limited risk effects of an actively hunting large carnivore. *Ecology Letters* 16(8), pp. 1023–
647 1030.

648 Mills, M.G.L. and Mills, M.E.J. 2014. Cheetah cub survival revisited: a re-evaluation of the
649 role of predation, especially by lions, and implications for conservation. *Journal of Zoology*
650 292(2), pp. 136–141.

651 Moller, A.P. 2004. Protandry, sexual selection and climate change. *Global Change Biology*
652 10(12), pp. 2028–2035.

653 Morehouse, A.T. et al. 2016. Nature vs. nurture: evidence for social learning of conflict
654 behaviour in grizzly bears. *PLOS ONE* 11(11), p. e0165425.

655 Muhly, T.B. et al. 2011. Human activity helps prey win the predator-prey space race. *PLoS*
656 *ONE* 6(3), p. e17050.

657 Newey, S. et al. 2015. Limitations of recreational camera traps for wildlife management and
658 conservation research: A practitioner’s perspective. *Ambio* 44(S4), pp. 624–635.

659 Ng, S.J. et al. 2004. Use of highway undercrossings by wildlife in southern California.
660 *Biological Conservation* 115(3), pp. 499–507.

661 Ngama, S. et al. 2016. How bees deter elephants: beehive trials with forest elephants
662 (*Loxodonta africana cyclotis*) in Gabon. *PLOS ONE* 11(5), p. e0155690.

663 Nolet, B.A. et al. 2006. Foraging costs and accessibility as determinants of giving-up densities
664 in a swan-pondweed system. *Oikos* 112(2), pp. 353–362.

665 Norris, D. et al. 2010. Habitat patch size modulates terrestrial mammal activity patterns in
666 Amazonian forest fragments. *Journal of Mammalogy* 91(3), pp. 551–560.

667 Nowak, K. et al. 2014. Human observers impact habituated samango monkeys’ perceived
668 landscape of fear. *Behavioral Ecology* 25(5), pp. 1199–1204.

669 Nyakaana, S. et al. 2001. DNA evidence for elephant social behaviour breakdown in Queen
670 Elizabeth National Park, Uganda. *Animal Conservation* 4(3), pp. 231–237.

671 O’Brien, T.G. et al. 2010. The Wildlife Picture Index: monitoring top trophic levels. *Animal*
672 *Conservation* 13(4), pp. 335–343.

673 Ordiz, A. et al. 2013. Lasting behavioural responses of brown bears to experimental encounters
674 with humans. *Journal of Applied Ecology* 50(2), pp. 306–314.

675 Orrock, J.L. 2004. Rodent foraging is affected by indirect, but not by direct, cues of predation
676 risk. *Behavioral Ecology* 15(3), pp. 433–437.

677 Otani, T. 2001. Measuring fig foraging frequency of the Yakushima macaque by using
678 automatic cameras. *Ecological Research* 16(1), pp. 49–54.

679 Picman, J. and Schriml, L.M. 1994. A camera study of temporal patterns of nest predation in
680 different habitats. *The Wilson Bulletin*, pp. 456–465.

681 Pienkowski, M.W. 1979. Differences in habitat requirements and distribution patterns of
682 plovers and sandpipers as investigated by studies of feeding behaviour. *Verhandlungen der*
683 *Ornithologische Gesellschaft in Bayern* 23, pp. 105–124.

684 du Preez, B.D. et al. 2014. To bait or not to bait: a comparison of camera-trapping methods for
685 estimating leopard *Panthera pardus* density. *Biological Conservation* 176, pp. 153–161.

686 Robertson, B.A. and Hutto, R.L. 2006. A framework for understanding ecological traps and an
687 evaluation of existing evidence. *Ecology* 87(5), pp. 1075–1085.

688 Romero-Muñoz, A. et al. 2010. Temporal separation between jaguar and puma in the dry
689 forests of southern Bolivia. *Journal of Tropical Ecology* 26(03), pp. 303–311.

690 Rovero, F. et al. 2013. ‘ Which camera trap type and how many do I need?’ A review of camera
691 features and study designs for a range of wildlife research applications. *Hystrix, the Italian*
692 *Journal of Mammalogy* 24(2), pp. 148–156.

693 Rovero, F. and Zimmermann, F. 2016. *Camera Trapping for Wildlife Research*. Pelagic
694 Publishing Ltd.

695 Rowcliffe, J.M. et al. 2008. Estimating animal density using camera traps without the need for
696 individual recognition. *Journal of Applied Ecology* 45(4), pp. 1228–1236.

697 Rowcliffe, J.M. et al. 2014. Quantifying levels of animal activity using camera trap data.
698 *Methods in Ecology and Evolution* 5(11), pp. 1170–1179.

699 Rowcliffe, J.M. et al. 2016. Wildlife speed cameras: measuring animal travel speed and day
700 range using camera traps. *Remote Sensing in Ecology and Conservation* 2(2), pp. 84–94.

701 Rowell, T.E. 1966. Hierarchy in the organization of a captive baboon group. *Animal Behaviour*
702 14(4), pp. 430–443.

703 Rychlik, L. 2005. Overlap of temporal niches among four sympatric species of shrews. *Acta*
704 *Theriological* 50(2), pp. 175–188.

705 Schaller, G.B. 1967. *The Deer and the Tiger: A Study of Wildlife in India*. Chicago: University
706 of Chicago Press.

707 Severud, W.J. et al. 2011. Predator cues reduce American beaver use of foraging trails. *Human-*
708 *Wildlife Interactions* 5, pp. 296–305.

709 Shamoun-Baranes, J. et al. 2012. From sensor data to animal behaviour: an oystercatcher
710 example. *PLoS ONE* 7(5), p. e37997.

711 Steenweg, R. et al. 2016. Scaling up camera traps: monitoring the planet’s biodiversity with
712 networks of remote sensors. *Frontiers in Ecology and the Environment*.

713 Stewart, F.E.C. et al. 2016. Wolverine behavior varies spatially with anthropogenic footprint:
 714 implications for conservation and inferences about declines. *Ecology and Evolution* 6(5), pp.
 715 1493–1503.

716 Steyaert, S.M.J.G. et al. 2016. Human shields mediate sexual conflict in a top predator.
 717 *Proceedings of the Royal Society B: Biological Sciences* 283(1833), p. 20160906.

718 Sueur, C. et al. 2011. How can social network analysis improve the study of primate behavior?
 719 *American Journal of Primatology* 73(8), pp. 703–719.

720 Suraci, J.P. et al. 2016. A new Automated Behavioural Response system to integrate playback
 721 experiments into camera trap studies. *Methods in Ecology and Evolution*. Available at:
 722 <http://doi.wiley.com/10.1111/2041-210X.12711> [Accessed: 22 December 2016].

723 Suselbeek, L. et al. 2014. Food acquisition and predator avoidance in a Neotropical rodent.
 724 *Animal Behaviour* 88, pp. 41–48.

725 Swanson, A.M. et al. 2015. Snapshot Serengeti, high-frequency annotated camera trap images
 726 of 40 mammalian species in an African savanna. *Scientific Data* 2.

727 Sweitzer, R.A. and Furnas, B.J. 2016. Data from camera surveys identifying co-occurrence and
 728 occupancy linkages between fishers (*Pekania pennanti*), rodent prey, mesocarnivores, and
 729 larger predators in mixed-conifer forests. *Data in Brief* 6, pp. 783–792.

730 Tan, C.L. et al. 2013. Into the night: camera traps reveal nocturnal activity in a presumptive
 731 diurnal primate, *Rhinopithecus brelichi*. *Primates* 54(1), pp. 1–6.

732 Tobler, M.W. et al. 2008a. An evaluation of camera traps for inventorying large- and medium-
 733 sized terrestrial rainforest mammals. *Animal Conservation* 11(3), pp. 169–178.

734 Tobler, M.W. et al. 2008b. Further notes on the analysis of mammal inventory data collected
 735 with camera traps. *Animal Conservation* 11(3), pp. 187–189.

736 Villette, P. et al. 2017. Evaluating camera traps as an alternative to live trapping for estimating
 737 the density of snowshoe hares (*Lepus americanus*) and red squirrels (*Tamiasciurus*
 738 *hudsonicus*). *European Journal of Wildlife Research* 63(7). Available at:
 739 <http://link.springer.com/10.1007/s10344-016-1064-3> [Accessed: 18 December 2016].

740 Voelkl, B. and Kasper, C. 2009. Social structure of primate interaction networks facilitates the
 741 emergence of cooperation. *Biology Letters* 5(4), pp. 462–464.

742 Vogt, K. et al. 2014. Scent-marking behaviour and social dynamics in a wild population of
 743 Eurasian lynx *Lynx lynx*. *Behavioural Processes* 106, pp. 98–106.

744 Wang, Y. et al. 2015. Mesopredator spatial and temporal responses to large predators and
 745 human development in the Santa Cruz Mountains of California. *Biological Conservation* 190,
 746 pp. 23–33.

747 Ware, J.V. et al. 2015. Validation of mercury tip-switch and accelerometer activity sensors for
 748 identifying resting and active behavior in bears. *Ursus* 26(2), pp. 86–96.

749 Weerakoon, M.K. et al. 2014. Can camera traps be used to estimate small mammal population
750 size. In: Fleming, P. et al. eds. *Camera trapping: wildlife management and research*. CSIRO
751 Publishing, pp. 307–316.

752 Wey, T. et al. 2008. Social network analysis of animal behaviour: a promising tool for the study
753 of sociality. *Animal Behaviour* 75(2), pp. 333–344.

754 Whitehead, H. 2008. *Analyzing Animal Societies: Quantitative Methods for Vertebrate Social*
755 *Analysis*. University of Chicago Press.

756 Wilson, R.P. et al. 1986. Recording devices on free-ranging marine animals: does measurement
757 affect foraging performance? *Ecology* 67(4), pp. 1091–1093.

758 Wolf, J.B.W. et al. 2007. Social structure in a colonial mammal: unravelling hidden structural
759 layers and their foundations by network analysis. *Animal Behaviour* 74(5), pp. 1293–1302.

760 Wright, J.T. et al. 2010. Native species behaviour mitigates the impact of habitat-forming
761 invasive seaweed. *Oecologia* 163(2), pp. 527–534.

762 Zheng, X. et al. 2016. Individual identification of wild giant pandas from camera trap photos -
763 a systematic and hierarchical approach. *Journal of Zoology* 300, pp. 247–256.

764

Table 1. Potential advantages and disadvantages of three conventional methods commonly used to collect animal behavioural data. These are not necessarily contextual constants. For example, GPS accuracy is affected by vegetation density. Similarly, activity sensors may return detailed or simplistic data, depending on the device used. VHF = Radio telemetry tags; GPS = Global Positioning System tags; ACC = activity sensors; CT = camera traps.

Advantages	Method			
	VHF	GPS	ACC	CT
Allows independent data verification			✓	✓
Collection of biometric data during deployment	✓	✓	✓	
Combined analysis of movement and trait-based data	✓ ^{1,2}	✓	✓	✓
Detailed data ^{2,3,4*}		✓	✓	✓
Habitat associations	✓	✓		✓
Identification of specific behaviours			✓*	✓
Landscape-scale	✓	✓		✓
Low cost			✓	✓*
Low survey effort		✓*	✓*	✓*
Multi-taxa surveys				✓
Range analyses	✓	✓		✓
Disadvantages				
Bias from handling focal animal(s) ^{5,8}	✓	✓	✓	
Disturbance effects				✓*
Expensive	✓	✓		✓*
Limited sample size	✓	✓	✓	
Negative impacts on focal animal(s) during backpack/collar deployment ⁷	✓	✓	✓	
Requires ground-truthing to avoid inferential error ^{4,5,6}			✓	
Simplistic data*	✓	✓	✓ ⁹	✓
Stationary				✓
Technological failure	✓	✓	✓	✓
Triangulation/location error ⁵	✓	✓		

* Device, environment and/or species-dependent

¹ Grignolio et al. (2004)

² Lewis et al. (2002)

³ Bouten et al. (2013)

⁴ Shamoun-Baranes et al. (2012)

⁵ Bridges and Noss (2011)

⁶ Ware et al. (2015)

⁷ Barron et al. (2010)

⁸ Wilson et al. (1986)

⁹ Coulombe et al. (2006)

Table 2. Examples of behavioural observations of wildlife via camera trapping. Species are ordered chronologically following the date of corresponding references.

Behaviour	Species	References
Active period	Agouti (<i>Dasyprocta punctata</i>) and ocelot (<i>Leopardus pardalis</i>)	Suselbeek et al. 2014
	Guizhou snub-nosed monkey (<i>Rhinopithecus brelichi</i>)	Claridge et al. 2004
	Spotted-tailed quoll (<i>Dasyurus maculatus</i>)	Tan et al. 2013
Antipredator responses	Bush rat (<i>Rattus fuscipes</i>)	Carthey and Banks 2016
Bathing/wallowing	Giant anteater (<i>Myrmecophaga tridactyla</i>)	Emmons et al. 2004
Crossing roads	Bare-nosed wombats (<i>Vombatus ursinus</i>)	Crook et al. 2013
Daily activity	Clouded leopard (<i>Neofelis nebulosa</i>), golden cat (<i>Catopuma temminckii</i>), and 4 other felids	Azlan and Sharma 2006
	Tayra (<i>Eira barbara</i>)	Delgado-V. et al. 2011
	Giant otter (<i>Pteronura brasiliensis</i>)	Leuchtenberger et al. 2014
	12 terrestrial mammal species	Rowcliffe et al. 2014
Denning	American black bear (<i>Ursus americanus</i>)	Bridges et al. 2004
Foraging	Yakushima macaque (<i>Macaca fuscata yakui</i>)	Otani 2001
	Tayra (<i>Eira barbara</i>)	Delgado-V. et al. 2011
Migration	Bald eagle (<i>Haliaeetus leucocephalus</i>), black vulture (<i>Coragyps atratus</i>) and 5 other birds of prey	Jachowski et al. 2015
Nest predation	Predators exploiting quail (<i>Coturnix coturnix</i>) eggs	Picman and Schriml 1994
Phenological changes	Elk (<i>Cervus elaphus</i>)	Brodie et al. 2012
Positional behaviour	Bare-tailed woolly opossum (<i>Caluromys philander</i>)	Dalloz et al. 2012
Resource partitioning	Cape fox (<i>Vulpes chama</i>), caracal (<i>Caracal caracal</i>), honey badger (<i>Mellivora capensis</i>) and 9 other carnivores	Edwards et al. 2015
Response to human-animal conflict	Tiger (<i>Panthera tigris</i>) and associated prey species	Johnson et al. 2006
Scent marking	Tayra (<i>Eira barbara</i>)	Delgado-V. et al. 2011
	Eurasian lynx (<i>Lynx lynx</i>)	Vogt et al. 2014
Social behaviour	Blonde capuchin (<i>Sapajus flavius</i>)	Bezerra et al. 2014
	Giant otter (<i>Pteronura brasiliensis</i>)	Leuchtenberger et al. 2014
Temporal avoidance	Jaguar (<i>Panthera onca</i>) and puma (<i>Puma concolor</i>)	Romero-Muñoz et al. 2010
Travel speed	12 terrestrial mammal species	Rowcliffe et al. 2016
Waterhole use	15 species of ungulates, 5 birds, 3 mega-herbivores, 2 primates and 5 carnivores	Hayward and Hayward 2012




Figure 1. Examples of animal behaviour captured by camera traps: **a)** Scent marking by an American black bear (*Ursus americanus*); **b)** intraspecific competition in moose (*Alces alces*); **c)** interspecific interactions between a European hare (*Lepus europaeus*; anti-predator response), a common buzzard (*Buteo buteo*; avoidance and attempted predation) and a hooded crow (*Corvus cornix*; anti-predator behaviour) captured on video (available at [10.6084/m9.figshare.4508369](https://doi.org/10.6084/m9.figshare.4508369)); **d)** predation of a European rabbit (*Oryctolagus cuniculus*) by a red fox (*Vulpes vulpes*); **e)** investigation of a squirrel feeding station by a pine marten (*Martes martes*); **f)** nut caching by a grey squirrel (*Sciurus carolinensis*). Images provided by A.C. Burton (a, b), A. Caravaggi (c, d) and C.M.V. Finlay (e, f).